

GUJARAT TECHNOLOGICAL UNIVERSITY**BE- Vth SEMESTER-EXAMINATION – MAY/JUNE - 2012****Subject code: 151002****Date: 04/06/2012****Subject Name: Engineering Electromagnetics****Time: 02:30 pm – 05:00 pm****Total Marks: 70****Instructions:**

1. Attempt all questions.
2. Make suitable assumptions wherever necessary.
3. Figures to the right indicate full marks.

Q.1 (a) Given points are E (2, 5, 1), F (-1, 4, -2) and G (3, -2, 4). **07**

Find: (i) A unit vector directed from E towards F.

(ii) The angle between $\mathbf{R}_{EF}$ and $\mathbf{R}_{EG}$.(iii) The scalar projection $\mathbf{R}_{EF}$ and $\mathbf{R}_{EG}$.

(b) (i) Find the volume defined by $4 < \rho < 6$, $30^\circ < \phi < 60^\circ$, $2 < z < 5$. (ii) What is the length of the longest straight line that lies entirely within the volume? (iii) Find the total area of the surface. **07**

Q.2 (a) Find Electric field intensity due to infinite line with uniform line charge density ρ_L on z-axis **07**

(b) Find E at P (1, 5, 2) in free space if a point charge of $6 \mu\text{C}$ is located at Q(0, 0, 1), a uniform line charge $\rho_L = 180 \text{ nC/m}$ lies along the z axis and a uniform sheet charge 25 nC/m^2 lies in the plane $z = -1$ **07**

OR

(b) Given the field $\bar{\mathbf{D}} = 6\rho \sin(\phi/2) \bar{\mathbf{a}}_\rho + 1.5\rho \cos(\phi/2) \bar{\mathbf{a}}_\phi \text{ C/m}^2$. **07**
Evaluate both sides of the divergence theorem for the region bounded by $\rho = 2$, $0 < \phi < 180^\circ$, $0 < z < 5$.

Q.3 (a) Derive equation to find Energy stored in the field of a system of charges **07**

(b) The region $y < 0$ contains a dielectric material for which $\epsilon_{R1} = 2.5$, while the region $y > 0$ is characterized by $\epsilon_{R2} = 4$. Let $\bar{\mathbf{E}}_1 = -30 \bar{\mathbf{a}}_x + 50 \bar{\mathbf{a}}_y + 70 \bar{\mathbf{a}}_z \text{ V/m}$ Find (i) D_{N2} (ii) D_{t2} (iii) $\bar{\mathbf{D}}_2$ (iv) θ_2 **07**


OR**Q.3 (a)** Find Capacitance between two concentric spheres using Laplace Equation. **07**

(b) Given a point of $200\epsilon_0 \text{ C}$ at (3, -1, 2), a line charge of $40\epsilon_0 \text{ C/m}$ on the x-axis and a surface charge of $8\epsilon_0 \text{ C/m}^2$ on the plane $x = -3$, all in free space. Find the potential at P (5, 6, 7) if $V = 0$ at Q (0, 0, 1). **07**

- Q.4** (a) Define the term “curl”. Also explain the point form of “Ampere’s circuital law”. **07**
 (b) Write the statement of Stoke’s theorem. **07**
 Evaluate both sides of Stoke’s theorem for the field $\vec{H} = 6xy \vec{a}_x - 3y^2 \vec{a}_y$ A / m and the rectangular path around the region, $2 \leq x \leq 5$, $-1 \leq y \leq 1$, $z = 0$. Let the positive direction of dS be $\vec{a}_z$.

OR

- Q.4** (a) Prove that for a differential current loop which carries current I in a given magnetic field, the torque on that loop is given by $d\vec{T} = d\vec{m} \times \vec{B}$. **07**
 (b) As shown in the figure below, the finite length current element is on the z – axis. Using Bio–savart law show that $\vec{H} = I / 4\pi\rho (\sin\alpha_2 - \sin\alpha_1) \vec{a}_\phi$. **07**



- Q.5** (a) Derive Maxwell’s four equations in point form. **07**
 (b) Explain Skin effect in detail. A steel pipe is constructed of a material for which $\mu_r = 200$ and $\sigma = 5 \times 10^6$ mho/m. The outer and inner radii 8 and 6 mm respectively and the length is 80 m. If the total current carried by the pipe is $2\cos 10^4 \pi t$ A, find (i) the skin depth (ii) the effective resistance **07**

OR

- Q.5** (a) Derive Maxwell’s four equations in integral form. **07**
 (b) A lossy dielectric is characterized by $\epsilon_r = 2.5$, $\mu_r = 4$, and $\sigma = 10^{-3}$ mho/m at 10MHz. Let $\vec{E}_s = 20 e^{-yz} \vec{a}_x$ V/m and find (i) α (ii) η (iii) $\vec{H}_s$ **07**
