

GUJARAT TECHNOLOGICAL UNIVERSITY**M. E. - SEMESTER – II • EXAMINATION – WINTER • 2014****Subject code: 1722309****Date: 08-12-2014****Subject Name: Numerical Methods****Time: 02:30 pm - 05:00 pm****Total Marks: 70****Instructions:**

1. Attempt all questions.
2. Make suitable assumptions wherever necessary.
3. Figures to the right indicate full marks.

- Q.1 (a)** Write algorithm of Newton Raphson Method. Also, explain its pitfalls. **07**
- (b)** Use Bisection method to determine drag coefficient c for a parachutist after free falling, using the nonlinear equation $f(c) = \left(\frac{667.38}{c} \right) (1 - e^{-(0.146843c)}) - 40 = 0$. Taking initial guesses (14.5 15) and stopping criteria 0.5%. **07**

- Q.2 (a)** Give geometrical interpretation of False Position method and use it to find the real root of $f(x) = xe^x - 2 = 0$ which lies between 0.8 and 0.9 correct to three decimal places. **07**
- (b)** What is mathematical modeling? Develop a model to find the velocity v of a freely falling parachutist as a function of time assuming that the air resistance is linearly proportional to v , How can you solve the problem numerically? **07**

OR

- (b)** Use following table to compute value of the Bessel function $J_0(x)$ for **07**
- 1) $x=1.72$ using Newton's forward difference interpolation formula and
 - 2) $x=1.93$ using Newton's backward difference interpolation formula.

x	1.7	1.8	1.9	2.0
$f(x)$	0.397 9849	0.339 9864	0.281 8186	0.223 8908

- Q.3 (a)** Describe Algorithm for the Gauss- Seidel Method. **07**
- (b)** 1) What is partial pivoting? Why do we use it? **07**
- 2) Use Gauss Elimination method to solve the following system of equations.
- $$3x + 2y + z = 10$$
- $$2x + 3y + 2z = 14$$
- $$x + 2y + 3z = 14$$

OR

- Q.3 (a)** For a spring mass system, following equations are obtained. Use Gauss Jordan method to find displacements x_1, x_2, x_3 **07**
- $$30x_1 - 20x_2 = 19.6$$
- $$-20x_1 + 30x_2 - 10x_3 = 29.4$$
- $$-10x_2 + 10x_3 = 24.5$$
- (b)** The following system of equations was generated by applying Kirchhoff's law to an electric circuit. Use Gauss Seidel method upto three iterations to determine the currents I_1, I_2 and I_3 . **07**
- $$3I_1 - 0.1I_2 - 0.2I_3 = 7.85$$
- $$0.1I_1 + 7I_2 - 0.3I_3 = -19.3$$
- $$0.3I_1 - 0.2I_2 + 10I_3 = 71.4$$

- Q.4 (a)** Derive normal equations for polynomial regression to fit a second degree polynomial for the given data points $(x_i, y_i), i = 1, 2, \dots, n$ using least square method. **07**

- (b)** Fit first order Spline to the data given in the following table. Also, evaluate the function at $x = 5$. **07**

x	3	4.5	7	9
$f(x)$	2.5	1.0	2.5	0.5

OR

- Q.4 (a)** What is difference between the two methods of curve fitting, namely Regression and interpolation? **07**

The voltage v across a capacitor at time t second is given by the following table:

Use the method of least square to fit a curve $v = ae^{kt}$ to this data.

t	0	2	4	6	8
v	150	63	28	12	5.6

- (b)** The following data was taken from an experiment. Use linear regression to fit a straight line for data given in the following table. Also, find the standard error of the estimate. **07**

X	1	2	3	4	5	6	7
Y	0.5	2.5	2.0	4.0	3.5	6.0	5.5

- Q.5 (a)** 1) Why do we need methods of Numerical Integration? **02**

2) A wind force distributed against the side of a skyscraper is measured as **05**

Height L (m)	0	30	60	90	120	150	180	210	240
Force F (N/m)	0	350	1000	1500	2600	3000	3300	3500	3600

Calculate the net force using Simpson's 1/3 rule.

- (b)** Use Runge-Kutta fourth order method to calculate $y(0.2)$ given $\frac{dy}{dx} = x + y$, $y(0) = 1$, taking $h = 0.1$. **07**

OR

- Q.5 (a)** Use Heun's method to integrate $y' = \frac{2y}{x}$, $y(1) = 2$ to estimate $y(1.5)$, taking step size $h = 0.25$ and only one iteration in the corrector step. **07**

- (b)** A steady state heat balance for a 10-m rod can be represented as **07**
- $$\frac{d^2T}{dx^2} + h'(T_a - T) = 0$$
- with $h' = 0.01 \text{ m}^{-2}$, $T_a = 20^\circ \text{C}$, $T(0) = 40$ and $T(10) = 200$
- Use the finite difference approach with $\Delta x = 2 \text{ m}$ to solve the boundary value problem.

□□□□□□□□□□